



EUROPEAN
HEMATOLOGY
ASSOCIATION

EHA25 VIRTUAL

25th Congress of the European Hematology Association

VIRTUAL EDITION



EUROPEAN
HEMATOLOGY
ASSOCIATION

EHA25 VIRTUAL

Interim Analysis From the OPTIC Trial: A Dose-Ranging Study of 3 Starting Doses of Ponatinib

Jorge Cortes,¹ Elza Lomaia,² Anna Turkina,³ Beatriz Moiraghi,⁴ Maria Undurraga Sutton,⁵ Carolina Pavlovsky,⁶ Christine Rojas,⁷ Charles Chuah,⁸ Christopher Kevin Arthur,⁹ Jane Apperley,¹⁰ Dong-Wook Kim,¹¹ Andreas Hochhaus,¹² Philippe Rousselot,¹³ Gianantonio Rosti,¹⁴ Michael Mauro,¹⁵ Jeffrey Lipton,¹⁶ Daniel Naranjo,¹⁷ Guohui Liu,¹⁷ Shouryadeep Srivastava,¹⁷ Michael Deininger¹⁸

¹Georgia Cancer Center, Augusta, GA, USA; ²Almazov National Medical Research Centre, Moscow, Russia; ³Russian Academy of Medical Sciences, Moscow, Russia; ⁴Hospital Jose Maria Ramos Mejia, Buenos Aires, Argentina; ⁵Hospital del Salvador, Santiago, Chile; ⁶Fundaleu, Buenos Aires, Argentina; ⁷Centro de Investigaciones Clinicas Vina del Mar, Valparaíso, Chile; ⁸Singapore General Hospital, Duke-NUS Medical School, Singapore; ⁹Royal North Shore Hospital, St. Leonards, Australia; ¹⁰Imperial College Healthcare NHS Trust, London, United Kingdom; ¹¹Catholic University of Korea – Seoul, Seoul, South Korea; ¹²Universitätsklinikum Jena, Jena, Germany; ¹³Hospital Mignot University de Versailles Saint-Quentin-en-Yvelines, Paris, France; ¹⁴University of Bologna, Bologna, Italy; ¹⁵Memorial Sloan Kettering, New York, NY, USA; ¹⁶University of Toronto, Toronto, Ontario, Canada; ¹⁷Millennium Pharmaceuticals, Inc., Cambridge, MA, USA, a wholly owned subsidiary of Takeda Pharmaceutical Company Limited; ¹⁸University of Utah, Huntsman Cancer Hospital, Salt Lake City, UT, USA


June 12, 2020

CML Clinical

Introduction

- Ponatinib is an orally active third-generation TKI designed to optimally inhibit all BCR-ABL1 single mutants^{1,2}
- Patients with resistant and intolerant CP-CML with substantial prior treatment demonstrated deep, lasting responses to ponatinib in the pivotal PACE trial¹
- A post hoc analysis modeling the data from PACE suggested a relationship between dose and both adverse events (AEs) and response rates³
- Lower ponatinib doses may mitigate AOE risk while maintaining efficacy³
- At this interim analysis, we present 21-month results (data cutoff: 20 July 2019) from the OPTIC trial, which evaluates the safety and efficacy of ponatinib over a range of 3 starting doses (45, 30, or 15 mg/d)

OPTIC (Optimizing Ponatinib Treatment In CP-CML): Ongoing, Multicenter, Randomized Phase 2 Trial



^a Dose reductions due to AEs were permitted

^b Escalation to the starting dose allowed for patients who lose their response following dose reduction; no dose escalation allowed beyond starting dose

AOE, arterial occlusive event; CCyR, complete cytogenetic response; CML, chronic myeloid leukemia; CV, cardiovascular; IA, interim analysis; IS, International Scale; MCyR, major cytogenetic response; MMR, major molecular response; VTE, venous thromboembolism



EUROPEAN
HEMATOLOGY
ASSOCIATION


EHA25 VIRTUAL

Patient Demographics and Baseline Disease Characteristics

Demographic/Disease Characteristic		Cohort A 45 mg (n=94)	Cohort B 30 mg (n=94)	Cohort C 15 mg (n=94)
Age, y	Median (range)	46.0 (19–81)	50.5 (21–77)	49.0 (18–81)
Gender, n (%)	Male	50 (53.2)	38 (40.4)	53 (56.4)
ECOG PS score, n (%)	0 or 1	93 (98.9)	93 (98.9)	94 (100)
Time since diagnosis, y	Median (range)	5.5 (1–21)	4.9 (1–29)	5.7 (1–22)
Prior TKIs, n (%)	1	1 (1.1)	1 (1.1)	4 (4.3)
	2	43 (45.7)	37 (39.4)	42 (44.7)
	≥3	50 (53.2)	56 (59.6)	48 (51.0)
BCR-ABL1 mutation at baseline, n (%)	No mutation detected	51 (54.3)	58 (61.7)	54 (57.4)
	<i>T315I</i>	25 (26.6)	21 (22.3)	20 (21.3)
	1 mutation detected	31 (33.0)	29 (30.9)	33 (35.1)
	≥2 mutations detected	10 (10.6)	6 (6.4)	5 (5.4)
Reason prior therapy stopped, n (%)	Resistant	93 (98.9)	94 (100.0)	94 (100.0)
Best response to last prior TKI, n (%)	None/PD	29 (31)	27 (29)	18 (19)
	CHR	31 (33)	28 (30)	39 (41)
	MCyR	20 (21)	16 (17)	10 (11)
	CCyR	7 (7)	6 (6)	7 (7)
	MMR	2 (2)	6 (6)	7 (7)

CHR, complete hematologic response; PD, progressive disease; SD, standard deviation

Patient Disposition



By the IA (cutoff date of July 20, 2019), 77% (n/N=216/282) of patients in the OPTIC trial were evaluable for the primary endpoint

^a Includes patients who have reached 12 months' follow-up, those who had early termination (ie, not reached 12-month mark but discontinued treatment), and those who have the b2a2/b3a2 transcript type at baseline




EUROPEAN
HEMATOLOGY
ASSOCIATION

EHA25 VIRTUAL

Exposure and Follow-up by Ponatinib Starting Dose

Exposure		Cohort A 45 mg (n=94)	Cohort B 30 mg (n=94)	Cohort C 15 mg (n=94)
Duration of exposure, months	Median (range)	14 (0.1–43.6)	12 (0.1–46.2)	12 (0.3–51.6)
Duration of exposure, n (%)	<1 month	3 (3.2)	4 (4.3)	2 (2.1)
	1–<3 months	6 (6.4)	10 (10.6)	8 (8.5)
	3–<6 months	13 (13.8)	8 (8.5)	14 (14.9)
	6–<12 months	18 (19.1)	21 (22.3)	23 (24.5)
	≥12–<24 months	27 (28.7)	30 (31.9)	26 (27.7)
	≥24 months	27 (28.7)	21 (22.3)	21 (22.3)
Dose intensity, mg/d	Median (range)	30.0 (10.5–45.0)	24.3 (5.5–30.0)	15.0 (6.0–15.0)
Follow-up, months	Median (range)	21 (15.0–24.0)	20 (13.7–23.9)	21 (12.6–24.1)

BCR-ABL1^{IS} ≤1% and MMR at 12 Months



BCR-ABL1^{IS} ≤1% at Month 12, n (%) [95% CI]; BCR-ABL1^{IS} ≤0.1% (MMR) at Month 12, n (%). At this IA, the analysis for primary endpoint is descriptive and no statistical inference is to be made. CI, confidence interval. Patients who have not reached 12-month primary endpoint evaluable criteria were not included in the IA (Cohort A, n=18; Cohort B, n=20; Cohort C, n=22).

Maintenance of Response Upon Dose Reduction

	Cohort A, 45 mg (n=93)		Cohort B, 30 mg (n=93)	
	Number of Patients Who Achieved BCR-ABL1 ^{IS} ≤1%, n	Percent of Responders Who Maintained BCR-ABL1 ^{IS} ≤1%, n/n (%)	Number of Patients Who Achieved BCR-ABL1 ^{IS} ≤1%, n	Percent of Responders Who Maintained BCR-ABL1 ^{IS} ≤1% n/n (%)
TOTAL	48		33	
No dose reduction to 15 mg ^a	6		2	
Dose reduction to 15 mg <u>BEFORE</u> achieving BCR-ABL1 ^{IS} ≤1%	6	6/6 (100)	5	3/5 (60)
Dose reduction to 15 mg <u>AFTER</u> achieving BCR-ABL1 ^{IS} ≤1%	36	27/36 (75)	24	21/24 (88)
<90 days after dose reduction	13	4/13 (31)	10	7/10 (70)
≥90 days after dose reduction	23	23/23 (100)	14	14/14 (100)
≥180 days after dose reduction	19	19/19 (100)	13	13/13 (100)
≥360 days after dose reduction	12	12/12 (100)	8	8/8 (100)

- Molecular responses were maintained in all patients who were on the reduced dose for at least 90 days
- Patients who could not maintain their response on a reduced dose of 15 mg/d, can be identified in less than 90 days after dose reduction


^a Protocol deviation

Re-escalation^a of Dose in Patients Who Lost Response

	Cohort A (45 mg) (n=93)	Cohort B (30 mg) (n=93)	Cohort C (15 mg) (n=90)
Achieved $\leq 1\%$ BCR-ABL1 ^{IS} at any time, n	48	33	29
Loss of $\leq 1\%$ BCR-ABL1 ^{IS} at any time, n	9	5	2
Dose re-escalated after loss of response, n	9	4	0
Regained $\leq 1\%$ BCR-ABL1 ^{IS}			
Yes, n (%)	3 (33.3)	2 (50.0)	0
No, n (%)	6 (66.7)	2 (50.0)	0
Median (range) follow-up for patients with dose re-escalation after loss of response, days	134 (1–746)	392 (1–510)	–


^a Escalation was not allowed beyond the starting dose

Time to $\leq 1\%$ BCR-ABL1^{IS} Response by Ponatinib Starting Dose



	Number at risk							
	0	3	6	9	12	15	18	21
Cohort A	93	78	47	33	20	14	7	6
Cohort B	93	70	50	43	31	14	9	7
Cohort C	90	78	49	37	25	11	9	7

BCR-ABL1^{IS} Response at 12 Months^a by T315I Baseline Status




- In the 15-mg/d starting dose cohort, a lower proportion of patients with the T315I mutation at baseline achieved ≤1% BCR-ABL1^{IS} vs those without (8.3% vs 30.4%), suggesting that this starting dose is not as effective as the higher starting doses for patients with the T315I mutation


MMR is ≤0.1% BCR-ABL1^{IS}

^a Not all patients reached 12 months' follow-up at the time of the analysis

Progression-Free^a and Overall Survival^a



	Number at risk									
	0	6	12	18	24	30	36	42	48	54
Cohort A	94	72	54	38	26	11	3	1	0	0
Cohort B	95	74	53	23	18	9	3	2	0	0
Cohort C	94	69	46	30	20	7	3	1	1	0




	Number at risk									
	0	6	12	18	24	30	36	42	48	54
Cohort A	94	78	63	51	36	18	8	1	0	0
Cohort B	95	80	64	51	36	20	8	4	0	0
Cohort C	94	76	60	50	36	19	8	2	1	0

^a Not all patients reached 12 months' follow-up at the time of the analysis. PFS was analyzed according to the criteria in O'Brien et al, 2003 and included: death; development of accelerated-phase or blast-phase chronic myeloid leukemia; loss of CHR (in the absence of cytogenetic response) confirmed by development in complete blood counts at least 4 weeks apart; Loss of major cytogenetic response by bone marrow cytogenetic assessment; increasing white blood cell count in patient without CHR defined by doubling of white blood cell count to >20,000 on 2 occasions at least 4 weeks apart (after the first 4 weeks of therapy)


OS, overall survival; PFS, progression-free survival

Most Common Treatment-Emergent Adverse Events

Incidence of Most Common
Non-Hematologic TEAEs, %^{a,b}



Incidence of Most Common
Hematologic TEAEs, %^{a,c}



^a All patients (safety population); ^b AST, dry skin, abdominal pain, rash, hypertriglyceridemia, and pyrexia occurred in $\geq 10\%$ of the population within Cohort A or B, but not in the total population;

^c Thrombocytopenia includes platelet count decreased

ALT, alanine aminotransferase increased; AST, aspartate aminotransferase increased; TEAEs, treatment-emergent adverse events

TEAE Summary and Related Dose Modifications/ Discontinuations

TEAE Summary and Dose Modifications, n (%)	Cohort A 45 mg (n=94)	Cohort B 30 mg (n=94)	Cohort C 15 mg (n=94)
Any TEAE	97.9%	92.6%	91.5%
Grade ≥ 3 TEAEs	66.0%	56.4%	57.4%
SAEs	30.9%	23.4%	27.7%
Deaths on-study, n	2 ^a	--	2 ^b
TEAEs leading to treatment discontinuation, dose reduction, or dose interruption, n (%)	65 (69.1)	54 (57.4)	52 (55.3)
TEAE leading to dose reduction, n (%)	41 (43.6)	29 (30.9)	26 (27.7)
TEAE leading to dose interruption, n (%)	64 (68.1)	48 (51.1)	50 (53.2)
TEAE leading to treatment discontinuation, n (%)	17 (18.1)	14 (14.9)	13 (13.8)

^a Sudden death was the cause of death for both patients

^b Pneumonia was the cause of death for both patients

SAEs, serious adverse events

Pre-adjudication and Adjudicated AOE Rates and Dose Modifications

Treatment Emergent AOE, n (%)	Cohort A Ponatinib 45 mg		Cohort B Ponatinib 30 mg		Cohort C Ponatinib 15 mg	
	Pre-adjudication	Adjudicated	Pre-adjudication	Adjudicated	Pre-adjudication	Adjudicated
AOEs	8 (8.5)	5 (5.3)	4 (4.3)	4 (4.3)	2 (2.1)	1 (1.1)
Serious AOE	4 (4.3)	2 (2.1)	3 (3.2)	3 (3.2)	0	0
Grade ≥3 AOE	4 (4.3)	3 (3.2)	4 (4.3)	4 (4.3)	0	0
Dose modifications for AOE, n (%)	Cohort A Ponatinib 45 mg		Cohort B Ponatinib 30 mg		Cohort C Ponatinib 15 mg	
	Pre-adjudication	Adjudicated	Pre-adjudication	Adjudicated	Pre-adjudication	Adjudicated
Discontinuation	4 (4.3)	2 (2.1)	3 (3.2)	3 (3.2)	0	0
Reduction	1 (1.1)	0	1 (1.1)	1 (1.1)	0	0
Interruption	1 (1.1)	0	1 (1.1)	1 (1.1)	0	0

- Independent adjudication committee reviewed AOE using the ACC/AHA, FDA, and STCI definitions¹
- Blinded to dose, dose modification, and investigator causality opinion
- No AOE-related deaths reported
- Only 7 AOE (4 in A and 3 in B) led to treatment discontinuation
- Overall adjudication AOE rate of 3.5%

ACC/AHA, American College of Cardiology/American Heart Association; FDA, US Food and Drug Administration; STCI, Standardized Data Collection for Cardiovascular Trials Initiative


1. Hicks KA, et al. *J Am Coll Cardiol*. 2015;66(4):403-469.



EUROPEAN
HEMATOLOGY
ASSOCIATION

EHA25 VIRTUAL

Observed Ponatinib Dose Relationship With Response Rate and TE-AOE Incidence Rate



- The percentage of patients with $\leq 1\%$ BCR-ABL1^{IS} and TE-AOE rates increased with increasing ponatinib dose

^a Efficacy n's by cohort: 15 mg, n=90; 30 mg and 45 mg, n=93; ^b TE-AOE n's by cohort: all cohorts, n=94; TE-AOE, treatment-emergent arterial occlusive event

Conclusions

- OPTIC IA demonstrates benefit with ponatinib in all 3 dosing regimens in this resistant CP-CML population
- Maximum rates of $\leq 1\%$ BCR-ABL1^{IS} at 12 months were achieved in the 45-mg/d starting dose cohort, and responses were maintained with the dose reduction to 15 mg/d
- At this IA, response-based ponatinib dosing regimens resulted in a clinically manageable safety and AOE profile
- With a median follow-up of ≈ 21 months, IA from OPTIC demonstrates that optimal benefit:risk profile for ponatinib was achieved with a response-adjusted dosing regimen starting with 45-mg/d dose, followed by dose reduction to 15 mg/d upon achieving $\leq 1\%$ BCR-ABL1^{IS}
- Primary analysis will provide a refined understanding of the benefit:risk profile of 3 different starting doses of ponatinib

Acknowledgments

The authors would like to thank the patients, their families, and their caregivers; the OPTIC investigators and their team members at each study site; and colleagues from ARIAD Pharmaceuticals, Inc., Cambridge, MA, USA, a wholly owned subsidiary of Takeda Pharmaceutical Company Limited.

Professional medical writing assistance was provided Peloton Advantage, LLC, an OPEN Health company, Parsippany, NJ, USA, and funded by Millennium Pharmaceuticals, Inc., Cambridge, MA, USA, a wholly owned subsidiary of Takeda Pharmaceutical Company Limited.