

AACR

American Association
for Cancer Research®

**ANNUAL
MEETING**

2023

APRIL 14-19 • #AACR23



Development of a CDK2-Selective Small Molecule Inhibitor INCB123667 for the Treatment of *CCNE1*^{high} Breast Cancers

Saswati Chand, Michael Hansbury, Yvonne Lo, Michelle Kinder, Kathy Wang, Qian Wang Yao, Derek Zimmer, Katherine Drake, Pat Feldman, Justine Carl, Cynthia Timmers, Joshua Hummel, Susan Wee, Sunkyu Kim

Incyte Research Institute, Wilmington, DE

Disclosure Information

Saswati Chand


I have the following relevant financial relationships to disclose:

Employee of: Incyte Corporation

Stockholder in: Incyte Corporation

Development of Resistance to CDK4/6 Inhibitors Is Common in Advanced Breast Cancer

- Primary resistance: About 15% of patients treated with CDK4/6 inhibitor (CDK4/6i) + aromatase inhibitor, and up to 30% of those treated with CDK4/6i + fulvestrant, will develop recurrent disease within 6 months
- Acquired resistance: Almost all patients will eventually develop progressive disease
- Multiple pathways are implicated in resistance
- *CCNE1* amplification and cyclin E1 overexpression are
 - Predictive for resistance to CDK4/6i
 - Associated with poor clinical outcomes




Reprinted from Álvarez-Fernández M, Malumbres M. *Cancer Cell*. 2020;37:514-529. Copyright © 2020 with permission from Elsevier Inc.

High *CCNE1* Expression Is Predictive for Resistance to CDK4/6i in HR⁺HER2⁻ Breast Cancer


High *CCNE1* mRNA expression is associated with primary resistance to CDK4/6i palbociclib (PALOMA-3 trial)

A




- Low *CCNE1* expression, PAL + FUL (n=103; mPFS=14.1 months)
- Low *CCNE1* expression, placebo + FUL (n=48; mPFS=4.8 months)
- High *CCNE1* expression, PAL + FUL (n=91; mPFS=7.6 months)
- High *CCNE1* expression, placebo + FUL (n=60; mPFS=4.0 months)

B




CCNE1 Amplification Is Associated With Poor Prognosis in Patients With TNBC

**TNBCs with *CCNE1* amplification
(copy number >6) correlate with poor OS**



<i>CCNE1</i> Amp	TCGA	METABRIC
TNBC	9.48%	5.66%


**TNBCs with high *CCNE1* expression
correlate with poor OS (cBioPortal)**



CCNE1 Amplification Has Synthetic Lethality With CDK2 Inhibition

- *CCNE1* amplification and cyclin E overexpression in cancer cells are predictive of CDK2 dependency as demonstrated by genetic knockdown studies
- CDK2 in complex with cyclin E regulates the G1/S transition and promotes DNA replication during the cell cycle
- Patients with primary or acquired *CCNE1* amplification and cyclin E overexpression may benefit from CDK2-targeted therapy


Role of CDK2/CCNE1 Complex in the Cell Cycle



Used with permission of EUREKA SCIENCE (FZC) from Treating Neurodegenerative Conditions Through the Understanding of Neuronal Apoptosis. D'Mello SR, et al. *Curr Drug Targets - CNS Neurol Disord*. 2005;4(1); permission conveyed through Copyright Clearance Center, Inc. RB, retinoblastoma protein.

CCNE1-Amplified or Overexpressed Cancer Cells Are Dependent on CDK2

Project Achilles data from the DepMap portal indicate most CDK2-dependent cancer cell lines harbor *CCNE1* amplification or overexpression across multiple lineages



Biochemical and Cellular Assays Demonstrate INCB123667 Is a Potent and Selective CDK2 Inhibitor


CDK Target	Biochemical Assay IC ₅₀ , μM	Cellular Assay IC ₅₀ , μM
CDK2	0.00087	0.053
CDK1	0.195	0.692
CDK4	0.046	0.873
CDK6	0.206	1.582
CDK7	0.355	>10,000
CDK9	3.676	5.273

IC₅₀, half-maximal inhibitory concentration.
Wee S, et al. *Eur J Cancer*. 2022;174:S79.

INCB123667 Blocks G1/S Transition and Induces Senescence in *CCNE1*^{high} Breast Cancer Cells

INCB123667 blocks G1/S transition in *CCNE1*^{high} cells

A




DMSO, dimethyl sulfoxide.

Wee S, et al. *Eur J Cancer*. 2022;174:S79.

INCB123667 induces senescence in *CCNE1*^{high} cells



B



Senescence demonstrated by beta-galactosidase (SA-βgal) staining

INCB123667 Induces Potent pRB and Growth Inhibition in Palbociclib-Resistant *CCNE1*^{high} Breast Cancer Cell Lines


INCB123667 inhibits pRB in *CCNE1*^{high} breast cancer cells in vitro



DMSO, dimethyl sulfoxide; pRB, phosphorylated retinoblastoma protein.


INCB123667 Inhibited Tumor Growth in *CCNE1*^{high} HCC1569 Xenograft Model In Vivo

INCB123667 inhibited tumor growth by 66.5% ($P<0.001$) vs a tumor growth inhibition (TGI) of 34% by palbociclib in an HR⁻/HER2⁺ epithelial breast cancer line HCC1569 xenograft model in NSG/SCID mice




RB1 Loss Is Both an Intrinsic and Acquired Mechanism of Resistance to CDK4/6 Inhibition

INCB123667 inhibits RB wild-type and RB-mutant *CCNE1*^{high} cell growth in vitro





Cyclin E1/CDK2 remains active in RB-mutant *CCNE1*^{high} tumor cells




INCB123667 Is Active in RB Wild-Type and RB-Mutant Cells

***CCNE1*^{high} RB-mutant cancer lines are dependent on CDK2 in pan-cancer analysis**

INCB123667 inhibits tumor growth in NCI-H526 RB-mutant in vivo tumor model



First-in-Human Phase 1b Trial of INCB123667 in Selected *CCNE1*^{high} Patients With Advanced Malignancies



*No selection for *CCNE1*^{high}.
BC, breast cancer; CDK4/6i, CDK4/6 inhibitor; GI, gastrointestinal; RDE, recommended dose for expansion.
ClinicalTrials.gov Identifier: NCT05238922.

Summary

- INCB123667 is a selective and potent CDK2 kinase inhibitor with subnanomolar inhibitory activity against human CDK2/cyclin E1
- INCB123667 selectively inhibited RB phosphorylation, blocked G1-S transition, and induced cell growth inhibition in *CCNE1*^{high} preclinical breast cancer cell lines in vitro
- INCB123667 exhibited significant single-agent activity in vivo in *CCNE1*^{high} breast cancer xenograft
- INCB123667 inhibited RB-mutant and RB wild-type *CCNE1*^{high} cancer cells in vitro and in vivo
- A phase 1 clinical trial of INCB123667 in patients with advanced malignancies including *CCNE1*^{high} TNBC and HR⁺HER2⁻ tumors post-CDK4/6i is ongoing (NCT05238922)

Acknowledgements

- Kristine Stump
- Kanishk Kapilashrami
- Guofeng Zhang
- Manti Guha
- Erin Crowgey
- Maryanne Covington
- Ricardo Macarron
- This study was supported by Incyte Corporation
- Research and development services were provided by Pharmaron (Waltham, MA), Crown Bioscience (San Diego, CA), and Champions Oncology (Hackensack, NJ)
- Editorial and graphics support was provided by Envision Pharma Group (Philadelphia, PA) and funded by Incyte Corporation