



**SOLVE
ON.**

Lilly and Incyte provide update on supplemental New Drug Application for baricitinib for the treatment of moderate to severe atopic dermatitis

July 16, 2021

INDIANAPOLIS, July 16, 2021 /PRNewswire/ -- Eli Lilly and Company (NYSE: LLY) and Incyte (NASDAQ:INCY) announced today that the U.S. Food and Drug Administration (FDA) will not meet the Prescription Drug User Fee Act (PDUFA) action date for the supplemental new drug application (sNDA) for baricitinib for the treatment of adults with moderate to severe atopic dermatitis (AD). The delay is related to the FDA's ongoing assessment of JAK inhibitors.

"We are confident in the efficacy and safety of baricitinib data supporting a favorable benefit-risk profile for the treatment of atopic dermatitis and look forward to continuing to work with the FDA during the remainder of the review process," said Ilya Yuffa, senior vice president and president of Lilly Bio-Medicines. "We are committed to bringing baricitinib to market to help meet the needs for people living with atopic dermatitis."

Baricitinib, an oral JAK inhibitor discovered by Incyte and licensed to Lilly, is approved for the treatment of adults with moderately to severely active rheumatoid arthritis (RA) in more than 75 countries. It is also approved in over 40 countries for the treatment of adults with moderate to severe AD who are candidates for systemic therapy and approved in Japan for the treatment of certain hospitalized patients with COVID-19. Baricitinib is being studied in alopecia areata (AA), systemic lupus erythematosus (SLE) and juvenile idiopathic arthritis (JIA).

This action does not affect Lilly's previously issued financial guidance for 2021.

Indication and Usage for OLUMIANT (baricitinib) tablets (in the United States) for RA patients

OLUMIANT® (baricitinib) 2-mg is indicated for the treatment of adult patients with moderately to severely active rheumatoid arthritis who have had an inadequate response to one or more tumor necrosis factor (TNF) antagonist therapies. **Limitation of Use:** Not recommended for use in combination with other JAK inhibitors, biologic disease-modifying antirheumatic drugs (DMARDs), or with potent immunosuppressants such as azathioprine and cyclosporine.

IMPORTANT SAFETY INFORMATION FOR OLUMIANT (baricitinib) tablets

WARNING: SERIOUS INFECTIONS, MALIGNANCY, AND THROMBOSIS

SERIOUS INFECTIONS: Patients treated with Olumiant are at risk for developing serious infections that may lead to hospitalization or death. Most patients who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. If a serious infection develops, interrupt Olumiant until the infection is controlled. Reported infections include:

- **Active tuberculosis (TB), which may present with pulmonary or extrapulmonary disease. Test patients for latent TB before initiating Olumiant and during therapy. If positive, start treatment for latent infection prior to Olumiant use.**
- **Invasive fungal infections, including candidiasis and pneumocystosis. Patients with invasive fungal infections may present with disseminated, rather than localized, disease.**
- **Bacterial, viral, and other infections due to opportunistic pathogens.**

Carefully consider the risks and benefits of Olumiant prior to initiating therapy in patients with chronic or recurrent infection.

Closely monitor patients for the development of signs and symptoms of infection during and after treatment with Olumiant including the possible development of TB in patients who tested negative for latent TB infection prior to initiating therapy.

MALIGNANCIES: Lymphoma and other malignancies have been observed in patients treated with Olumiant.

THROMBOSIS: Thrombosis, including deep venous thrombosis (DVT) and pulmonary embolism (PE), has been observed at an increased incidence in patients treated with Olumiant compared to placebo. In addition, there were cases of arterial thrombosis. Many of these adverse events were serious and some resulted in death. Patients with symptoms of thrombosis should be promptly evaluated.

WARNINGS AND PRECAUTIONS

SERIOUS INFECTIONS: The most common serious infections reported with Olumiant included pneumonia, herpes zoster and urinary tract infection. Among opportunistic infections, tuberculosis, multidermatomal herpes zoster, esophageal candidiasis, pneumocystosis, acute histoplasmosis, cryptococcosis, cytomegalovirus and BK virus were reported with Olumiant. Some patients have presented with disseminated rather than localized disease and were often taking concomitant immunosuppressants such as methotrexate or corticosteroids. Avoid Olumiant in patients with an active, serious infection, including localized infections. Consider the risks and benefits of treatment prior to initiating Olumiant in patients:

- with chronic or recurrent infection
- who have been exposed to TB
- with a history of a serious or an opportunistic infection
- who have resided or traveled in areas of endemic tuberculosis or endemic mycoses; or
- with underlying conditions that may predispose them to infection.

Closely monitor patients for infections during and after Olumiant treatment. Interrupt Olumiant if a patient develops a serious infection, an opportunistic infection, or sepsis. Do not resume Olumiant until the infection is controlled.

Tuberculosis – Before initiating Olumiant evaluate and test patients for latent or active infection and treat patients with latent TB with standard antimycobacterial therapy. Olumiant should not be given to patients with active TB. Consider anti-TB therapy prior to initiating Olumiant in patients with a history of latent or active TB in whom an adequate course of treatment cannot be confirmed, and for patients with a negative test for latent TB but who have risk factors for TB infection. Monitor patients for TB during Olumiant treatment.

Viral Reactivation – Viral reactivation, including cases of herpes virus reactivation (e.g., herpes zoster), were reported in clinical studies with Olumiant. If a patient develops herpes zoster, interrupt Olumiant treatment until the episode resolves.

The impact of Olumiant on chronic viral hepatitis reactivation is unknown. Screen for viral hepatitis in accordance with clinical guidelines before initiating Olumiant.

MALIGNANCY AND LYMPHOPROLIFERATIVE DISORDERS: Malignancies were observed in Olumiant clinical studies. Consider the risks and benefits of Olumiant prior to initiating therapy in patients with a known malignancy other than a successfully treated non-melanoma skin cancer (NMSC) or when considering continuing Olumiant in patients who develop a malignancy. NMSCs were reported in patients treated with Olumiant. Periodic skin examination is recommended for patients who are at increased risk for skin cancer.

THROMBOSIS : Thrombosis, including DVT and PE, has been observed at an increased incidence in Olumiant-treated patients compared to placebo. In addition, arterial thrombosis events in the extremities have been reported in clinical studies with Olumiant. Many of these adverse events were serious and some resulted in death. There was no clear relationship between platelet count elevations and thrombotic events. Use Olumiant with caution in patients who may be at increased risk of thrombosis. If clinical features of DVT/PE or arterial thrombosis occur, evaluate patients promptly and treat appropriately.

GASTROINTESTINAL PERFORATIONS: Gastrointestinal perforations have been reported in Olumiant clinical studies, although the role of JAK inhibition in these events is not known. Use Olumiant with caution in patients who may be at increased risk for gastrointestinal perforation (e.g., patients with a history of diverticulitis). Promptly evaluate patients who present with new onset abdominal symptoms for early identification of gastrointestinal perforation.

LABORATORY ABNORMALITIES:

Neutropenia – Olumiant treatment was associated with an increased incidence of neutropenia (absolute neutrophil count [ANC] <1000 cells/mm³) compared to placebo. Avoid initiation or interrupt Olumiant treatment in patients with an ANC <1000 cells/mm³. Evaluate at baseline and thereafter according to routine patient management.

Lymphopenia – Absolute lymphocyte count (ALC) <500 cells/mm³ were reported in Olumiant clinical trials. Lymphocyte counts less than the lower limit of normal were associated with infection in patients treated with Olumiant, but not placebo. Avoid initiation or interrupt Olumiant treatment in patients with an ALC <500 cells/mm³. Evaluate at baseline and thereafter according to routine patient management.

Anemia – Decreases in hemoglobin levels to <8 g/dL were reported in Olumiant clinical trials. Avoid initiation or interrupt Olumiant treatment in patients with hemoglobin <8 g/dL. Evaluate at baseline and thereafter according to routine patient management.

Liver Enzyme Elevations – Olumiant treatment was associated with increased incidence of liver enzyme elevation compared to placebo. Increases of ALT ≥5x upper limit of normal (ULN) and increases of AST ≥10x ULN were observed in patients in Olumiant clinical trials.

Evaluate at baseline and thereafter according to routine patient management. Promptly investigate the cause of liver enzyme elevation to identify potential cases of drug-induced liver injury. If increases in ALT or AST are observed and drug-induced liver injury is suspected, interrupt Olumiant until this diagnosis is excluded.

Lipid Elevations – Treatment with Olumiant was associated with increases in lipid parameters, including total cholesterol, low-density lipoprotein cholesterol and high-density lipoprotein cholesterol. Assess lipid parameters approximately 12 weeks following Olumiant initiation. Manage patients according to clinical guidelines for the management of hyperlipidemia.

VACCINATIONS: Avoid use of live vaccines with Olumiant. Update immunizations in agreement with current immunization guidelines prior to initiating Olumiant therapy.

HYPERSENSITIVITY: Reactions such as angioedema, urticaria, and rash that may reflect drug sensitivity have been observed in patients receiving Olumiant, including serious reactions. If a serious hypersensitivity reaction occurs, promptly discontinue Olumiant while evaluating the potential causes of the reaction.

ADVERSE REACTIONS

Most common adverse reactions include: upper respiratory tract infections (16.3%, 11.7%), nausea (2.7%, 1.6%), herpes simplex (0.8%, 0.7%) and herpes zoster (1.0%, 0.4%) for Olumiant 2 mg and placebo, respectively.

USE IN SPECIFIC POPULATIONS

PREGNANCY AND LACTATION: No information is available to support the use of Olumiant in pregnancy or lactation. Advise women not to breastfeed during treatment with Olumiant.

HEPATIC AND RENAL IMPAIRMENT: Olumiant is not recommended in patients with severe hepatic impairment or in patients with severe renal impairment.

Please click to access full [Prescribing Information](#), including **Boxed Warning about Serious infections, Malignancies, and Thrombosis, and Medication Guide** .

About OLUMIANT® (baricitinib)

OLUMIANT, a once-daily, oral JAK inhibitor was discovered by Incyte and licensed to Lilly. It is approved in the U.S. and more than 75 countries as a treatment for adults with moderate to severe rheumatoid arthritis. It is also approved in over 40 countries for the treatment of adults with moderate to severe AD who are candidates for systemic therapy and for the treatment of certain hospitalized patients with COVID-19 in Japan. The U.S. FDA-approved labeling for OLUMIANT includes a Boxed Warning for Serious Infections, Malignancy, and Thrombosis. See the full Prescribing Information [here](#).

In December 2009, Lilly and Incyte announced an exclusive worldwide license and collaboration agreement for the development and commercialization of baricitinib and certain follow-on compounds for patients with inflammatory and autoimmune diseases.

About Lilly in Dermatology

By following the science through uncharted territory, we continue Lilly's legacy of delivering innovative medicines that address unmet needs and have significant impacts on people's lives around the world. Skin-related diseases are more than skin deep. We understand the devastating impact this can have on people's lives. At Lilly, we are relentlessly pursuing a robust dermatology pipeline to provide innovative, patient-centered solutions so patients with skin-related diseases can aspire to live life without limitations.

About Eli Lilly and Company

Lilly is a global health care leader that unites caring with discovery to create medicines that make life better for people around the world. We were founded more than a century ago by a man committed to creating high-quality medicines that meet real needs, and today we remain true to that mission in all our work. Across the globe, Lilly employees work to discover and bring life-changing medicines to those who need them, improve the understanding and management of disease, and give back to communities through philanthropy and volunteerism. To learn more about Lilly, please visit us at lilly.com and lilly.com/newsroom.

About Incyte

Incyte is a Wilmington, Delaware-based, global biopharmaceutical company focused on finding solutions for serious unmet medical needs through the discovery, development and commercialization of proprietary therapeutics. For additional information on Incyte, please visit lincyte.com and follow [@Incyte](https://twitter.com/Incyte).

P-LLY

OLUMIANT® is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995) about OLUMIANT (baricitinib) as a treatment for patients with rheumatoid arthritis and a possible treatment for patients with atopic dermatitis and other conditions and reflects Lilly's and Incyte's current beliefs and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the process of research, development, and commercialization. Among other things, there can be no guarantee that planned or ongoing studies will be completed as planned, that future study results will be consistent with the results to date, and that OLUMIANT will receive additional regulatory approvals, or be commercially successful. For further discussion of these and other risks and uncertainties, see Lilly's and Incyte's most recent respective Form 10-K and Form 10-Q filings with the United States Securities and Exchange Commission. Except as required by law, Lilly and Incyte undertake no duty to update forward-looking statements to reflect events after the date of this release.

Refer to: Kristen Basu; basu_kristen_porter@lilly.com; +1-317-447-2199 (Lilly media)
Kevin Hern; hern_kevin_r@lilly.com; +1-317-277-1838 (Lilly investors)
Catalina Loveman; cloveman@lincyte.com; +1-302-498-6171 (Incyte media)
Christine Chiou; cchiou@lincyte.com; +1-302-274-4773 (Incyte investors)

SOURCE Eli Lilly and Company