

RE-MIND study: comparison of tafasitamab + lenalidomide (L-MIND) vs lenalidomide monotherapy (real-world data) in transplant-ineligible patients with relapsed/refractory diffuse large B-cell lymphoma

Pier Luigi Zinzani,¹ Thomas Rodgers,² Dario Marino,³ Maurizio Frezzato,⁴ Anna Maria Barbui,⁵ Claudia Castellino,⁶ Erika Meli,⁷ Annarita Conconi,⁸ Nicola Cascavilla,⁹ Federica Cavallo,¹⁰ Nathan H. Fowler,¹¹ Bruce Feinberg,¹² Sascha Tillmanns,¹³ Stephan Parche,¹³ Günter Fingerle-Rowson,¹³ Mark Winderlich,¹³ Sumeet Ambarkhane,¹³ Gilles Salles,¹⁴ Grzegorz Nowakowski¹⁵

¹Institute of Hematology L e A Seràgnoli, University of Bologna, Bologna, Italy; ²University of Rochester Medical Center, Rochester, NY, United States;

³Medical Oncology 1, Department of Oncology, Veneto Institute of Oncology IOV-IRCCS, Padua, Italy; ⁴Azienda ULSS 8 Berica, Vicenza, Italy;

⁵ASST Papa Giovanni XXIII, Bergamo, Italy; ⁶Hematology Division, Santa Croce and Carle Hospital, Cuneo, Italy;

⁷ASST Grande Ospedale Metropolitano Niguarda, Dipartimento di Ematologia e Oncologia, S.C. Ematologia, Milan, Italy;

⁸SSD Ematologia, Ospedale degli Infermi di Biella, Ponderano, Italy; ⁹Istituto di Ematologia, IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo, Italy;

¹⁰Divisione Ematologia dell'Università di Torino, A.O.U. Città della Salute e della Scienza di Torino, Turin, Italy;

¹¹Department of Lymphoma/Myeloma, The University of Texas MD Anderson Cancer Center, Houston, TX, United States;

¹²Cardinal Health Inc., Dublin, OH, United States; ¹³MorphoSys AG, Planegg, Germany;

¹⁴Hématologie, Hospices Civils de Lyon and Université de Lyon, Lyon, France; ¹⁵Division of Hematology, Mayo Clinic, Rochester, MN, United States.

Disclosures

About tafasitamab (MOR208)

Tafasitamab (MOR208) is an investigational humanized Fc-engineered monoclonal antibody directed against CD19. In 2010, MorphoSys licensed exclusive worldwide rights to develop and commercialize tafasitamab from Xencor, Inc. Tafasitamab incorporates an XmAb® engineered Fc domain, which is intended to lead to a significant potentiation of antibody-dependent cell-mediated cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP), thus aiming to improve a key mechanism of tumor cell killing. XmAb® is a trademark of Xencor, Inc. Tafasitamab is being developed by MorphoSys and Incyte under a collaboration and license agreement dated January 12, 2020, to further develop and commercialize MorphoSys' proprietary anti-CD19 antibody tafasitamab (MOR208) globally.

Funded by MorphoSys AG; RE-MIND ClinicalTrials.gov number NCT04150328. PLZ reports receiving honoraria from Verastem, Celltrion, Merck, Gilead, Servier, Eusapharma, Debiopharm, Janssen, Roche, Abbvie, Takeda, Morphosys, MEI Pharma, Kirin Kyowa, TG Therapeutics. TR reports receiving personal fees for consultancy from Targeted Oncology. MF reports receiving a travel grant from Celgene. AMB reports receiving honoraria and personal fees for consultancy from Jansen as well as a travel grant from Roche. CC reports receiving institutional research funding from MorphoSys. NHF reports receiving personal fees for consultancy from Roche, BMS, Gilead and MorphoSys and reports receiving institutional research funding from Roche, BMS and Gilead. GS reports receiving personal fees for consultancy from Epizyme; advisory board participation from Amgen; advisory board participation or symposia from MorphoSys, Amgen, Acerta, Autolus, BMS, Merck, AbbVie, Janssen, Novartis, Gilead/Kite, Servier, Celgene, Roche, Pfizer and Takeda outside the submitted work; and has patent issued (WO2012010561A1: characterization of another anti-CD19 monoclonal antibody, having ADCC function, and developed in collaboration with IDD-biotech; this antibody and the company has no relationship with the anti-CD19 antibody described in the current paper (tafasitamab) and has not been licensed to any third party. GN reports receiving personal fees for consultancy from Celgene, BMS, Seattle Genetics, Roche/Genentech, MorphoSys, Selvita, Curis and reports receiving research funding from Celgene, BMS, Roche/Genentech and MorphoSys. ST, SP, GFR, SA and MW are employees of MorphoSys AG, Planegg, Germany. All other authors declare no competing interests

RE-MIND: background


- **Patients with non-transplant-eligible R/R DLBCL have a poor prognosis, with few available treatment options¹**
- **Tafasitamab is an Fc-enhanced humanized monoclonal anti-CD19 antibody that mediates increased ADCC, ADCP, and direct tumor cytotoxicity^{2,3}**
- **Lenalidomide (LEN), an immunomodulatory drug, is active in patients with R/R DLBCL, albeit with modest activity⁴⁻⁷**
- **L-MIND (NCT02399085) is an ongoing, open-label, single-arm, Phase II study of tafasitamab + LEN in patients with R/R DLBCL who are ineligible for ASCT⁸**
 - In the primary analysis of L-MIND, the IRC-assessed ORR was 60% with a CR rate of 42.5% and a median DoR of 21.7 months⁸
- **RE-MIND, a global real-world data study of LEN monotherapy, established a matched comparative cohort for the L-MIND study to isolate the contribution of tafasitamab to the tafasitamab + LEN combination⁹**

- ADCC, antibody-dependent cellular cytotoxicity; ADCP, antibody-dependent cellular phagocytosis;
ASCT, autologous stem cell transplantation; CR, complete response; DLBCL, diffuse large B-cell lymphoma;
DoR, duration of response; IRC, independent review committee; LEN, lenalidomide; ORR, overall response rate;
R/R, relapsed/refractory.
1. Friedberg JW. Hematology Am Soc Hematol Educ Program 2011;2011:498–505
 2. Awan FT, et al. Blood 2010;115:1204–13
 3. Horton HM, et al. Cancer Res 2008;68:8049–57
 4. Witzig TE, et al. Ann Oncol 2011;22(7):1622–7
 5. Czuczman MS, et al. Clin Cancer Res 2017;23(15):4127–37
 6. Wiernik PH, et al. J Clin Oncol 2008;26(30):4952–7
 7. Broccoli A, et al. Oncologist 2019;24(9):1246–52
 8. Salles G, et al. ICML 2019 (Abstract 124)
 9. Nowakowski G, et al. ASCO 2020 (Abstract 8020)

RE-MIND: trial design

NCT04150328

- Inclusion/exclusion criteria were aligned with those of the L-MIND trial¹
- Health record data of 490 patients were retrospectively collected via eCRF
- A control cohort for L-MIND comprised 76 patients with a LEN starting dose of 25 mg/day and matching for nine baseline characteristics via 1:1 ePS
- Multiple predefined sensitivity analyses, including alternative cohort balancing methodology and applying multiple imputation for missing values, demonstrated the robustness of the comparison




*Age (<70 vs ≥70 years); Ann Arbor stage (I/II vs III/IV); refractoriness to last therapy line (yes vs no); number of prior lines of therapy (1 vs 2 or 3); history of primary refractoriness (yes vs no); prior ASCT (yes vs no); elevated LDH (LDH >ULN vs LDH ≤ULN); neutropenia (ANC <1.5 × 10⁹/L vs ANC ≥1.5 × 10⁹/L); anemia (Hb <10 g/dL vs Hb ≥10 g/dL). †With an assumed difference of 23% in ORR, the achieved power is 80% and minimal detectable statistical difference is 17% under Fisher's exact test for unpaired data.

ANC, absolute neutrophil count; eCRF, electronic case report form; ePS, estimated propensity score; Hb, hemoglobin; Hx, history; INV, investigator-assessed; LDH, lactate dehydrogenase; OS, overall survival; PFS, progression-free survival; ULN, upper limit of normal.

1. Salles G, et al. ICML 2019 (Abstract 124)

RE-MIND: patient disposition




I/E, inclusion or exclusion criteria; tx, treatment.

EHA, June 11–14, 2020. Abstract: S238

Nowakowski G, et al. ASCO 2020 (Abstract 8020)

RE-MIND: baseline characteristics used for cohort balancing



Baseline characteristics were well balanced across the lenalidomide monotherapy and combination cohorts after the matching procedure

*SMD is defined as the ratio of the difference of proportions of a baseline characteristic to the standard deviation of the pooled difference. This standardization allows for comparison of the relative balance achieved across different baseline characteristics occurring in a low or high proportion.

SMD, standardized mean difference.

Nowakowski G, et al. ASCO 2020 (Abstract 8020)

EHA, June 11–14, 2020. Abstract: S238

RE-MIND: comparison of cohort characteristics

	Tafasitamab + LEN n=76	LEN monotherapy n=76
Median lenalidomide dose intensity,* mg/d (range)	17.6 (6.9–25.0)	19.0 (7.3–25.0)
Median follow-up for OS, [†] months	21.5	20.9
Median time to first post-baseline response assessment, months	1.9	3.1
Median frequency of response assessments, months	2.1	3.2
Imaging modality (CT or PET/CT), %	96	82

Cohort characteristics were comparable

*Cumulative dose divided by exposure time; [†]Reverse Kaplan–Meier analysis.


CT, computerized tomography; PET, positron emission tomography.

EHA, June 11–14, 2020. Abstract: S238

Nowakowski G, et al. ASCO 2020 (Abstract 8020)


RE-MIND: best overall response rate (primary endpoint)

Investigator-assessed




**The primary endpoint was met:
a significantly better ORR in the combination cohort versus the lenalidomide monotherapy cohort was shown**

RE-MIND: PFS and OS (secondary endpoints)



Significantly improved clinical efficacy (PFS and OS) was demonstrated for tafasitamab + lenalidomide compared with lenalidomide monotherapy

RE-MIND: sensitivity analyses for ORR



Sensitivity analyses, including achieving a higher degree of balance, utilizing alternative cohort balancing approaches and applying a multiple imputation technique, confirmed the primary analysis results

*Balancing weight approaches use weights based on the ePS to create a sample in which the distribution of measured baseline covariates is independent of treatment assignment, and estimate the average treatment effect in this population.

RE-MIND: summary

- **RE-MIND is a prospectively designed, high-quality real-world data study¹**
- **Outcomes observed for LEN monotherapy in RE-MIND compare well with published data from clinical trials^{2–5}**
- **A robust estimation of the additional treatment effect attributable to tafasitamab in the combination with LEN in transplant-ineligible R/R DLBCL was determined**
- **Tafasitamab + LEN provides significant and clinically meaningful improvements in ORR, CR rate, DoR, and OS compared with LEN monotherapy**
- **Effect sizes in all time-to-event endpoints and all sensitivity analyses support the primary endpoint results**
- **The study provides an important example of the utilization of real-world data to accelerate drug development**

RE-MIND demonstrates the significant contribution of tafasitamab to the efficacy observed in the combination with LEN, which represents a potential therapy option in R/R DLBCL

1. Nowakowski G, et al. ASCO 2020 (Abstract 8020)

2. Witzig TE, et al. Ann Oncol 2011;22(7):1622–7

3. Czuczman MS, et al. Clin Cancer Res 2017;23(15):4127–37

4. Wiernik PH, et al. J Clin Oncol 2008;26(30):4952–7

5. Broccoli A, et al. Oncologist 2019;24(9):1246–52